

## EPA-Approved Crypton Disinfectant & Deodorizer for Crypton Fabric

- **EPA-registered hospital-grade disinfectant**
  - Disinfects hard surfaces
  - Disinfects Crypton Barrier Fabrics
  - Safe for most other fabrics
- **Eliminates 99.9% of germs**
- **No-rinse formula**
- **One-step cleaner, deodorizer and disinfectant**
- **For use around children and pets**
- **Germicidal and virucidal performance with a fresh, clean scent**
- **Fights cold and flu**
- **Kills mold and mildew**
- **Deodorizes textiles**



**For use in:** Hospitals, nursing homes, medical and dental centers, patient care and recovery rooms, emergency rooms, labs, day care centers, nurseries, retirement facilities, restaurants, bars, kitchens, cafeterias, hotels, motels, dormitories, veterinary clinics, kennels, zoos, pet shops, cruise lines, airline terminals, trains, busses, trucks, public transportation, campgrounds, playgrounds, health clubs, spas, museums, theaters, banks, libraries, your home and any other place germs hide!

	Disinfects Crypton Fabric	Disinfects Hard Surfaces	Cleans and Disinfects in One Step	Low pH	Usable Around Pets	Kills Mold and Mildew	Eliminates Odor-causing Bacteria
Crypton Care Disinfectant & Deodorizer	•	•	•	•	•	•	•
Sanitizers				•			
Aerosol Disinfectants		•	•			•	•
Fragranced Fresheners							

## Crypton Disinfectant & Deodorizer

EPA-registered, one-step disinfectant cleaner effective against a broad spectrum of bacteria, viruses, mold, mildew and fungus, Crypton Disinfectant & Deodorizer disinfects hard, non-porous surfaces including Crypton barrier fabrics.

**Bactericidal (hospital disinfection):** Effective on hard inanimate surfaces; according to the AOAC Use Dilution Test method modified in the presence of 5% organic serum (850 ppm active). Treated surfaces must remain wet for 10 minutes.

Pseudomonas aeruginosa	Escherichia coli	Serratia marcescens
Salmonella enterica	Escherichia coli (antibiotic resistant)	Shigella dysenteriae
Staphylococcus aureus	Escherichia coli O157:H7	Shigella flexneri
Burkholderia cepacia	Klebsiella pneumoniae	Shigella sonnei
Campylobacter jejuni	Klebsiella pneumoniae (antibiotic resistant)	Methicillin Resistant
Corynebacterium ammoniagenes	Legionella pneumophila	Staphylococcus aureus (MRSA)
Enterobacter aerogenes	Listeria monocytogenes	Community Associates Methicillin Resistant
Enterobacter cloacae	Proteus mirabilis	Staphylococcus aureus (CA-MRSA)
Enterobacteriaceae with extended beta-lactamase resistance	Proteus vulgaris	Staphylococcus epidermidis (antibiotic resistant)
Enterococcus faecalis	Pseudomonas aeruginosa	Streptococcus pyogenes
Enterococcus faecium vancomycin resistant (VRE)	Salmonella typhi	Vibrio cholera

**Virucidal:** Effective on hard, non-porous environmental surfaces against the following viruses; Evaluated in the presence of 5% serum (850 ppm quat active) with a 10 minute contact time.

Avian Influenza A Virus (H5N1)	Herpes Simplex Type 2	Porcine Respiratory & Reproductive (PRRSV) Strain NVSL
Avian Influenza Turkey/Wisconsin	Human Coronavirus	Porcine Rotavirus
Canine Coronavirus	Human Immunodeficiency Virus Type 1 (HIV 1)	Pseudorabies Virus
Canine Distemper	Influenza A Virus (H1N1)	Respiratory Syncytial Virus
Hantavirus	Influenza A/Brazil Virus	Transmissible Gastroenteritis
Hepatitis B Virus	Infectious Bovine Rhinotracheitis Virus	Vaccinia Virus
Hepatitis C Virus	Newcastle Disease Virus	
Herpes Simplex Type 1		

**Non-food contact surface sanitizer:** Effective on hard non-porous non-food contact surfaces. Treated surfaces must remain wet for 5 minutes. Then wipe with sponge, mop or cloth or allow to air dry. Food contact surfaces must be rinsed.

Klebsiella pneumoniae  
Staphylococcus aureus

**Mold & mildew control:** Effective on hard, non-porous surfaces in controlling the growth of mold and mildew and the odors associated. Thoroughly wet all treated surfaces completely. Let air-dry. Repeat application weekly or when growth or odor reappears.

Aspergillus niger

**Fungicidal:** Effective against the following fungi on hard non-porous environmental surfaces; Evaluated at 2 ounces per gallon in the presence of 5% serum and 400 ppm hard water with a 10 minute contact time.

Candida albicans  
Dactylium dendroides  
Trichophyton mentagrophytes